



## Environment, Biodiversity & Soil Security (EBSS)

<http://jenvbs.journals.ekb.eg/>



### Nano-Selenium and its Interaction with other Nano-Nutrients in Soil under Stressful Plants: A Mini-Review



Hassan El-Ramady <sup>1</sup>, Naglaa Taha <sup>2</sup>, Tarek A. Shalaby <sup>3</sup>, Tamer Elsakhawy <sup>4</sup>, Alaa El-Dein Omara <sup>4</sup>, Jozsef Prokisch <sup>5</sup> and Yousry Bayoumi <sup>3,6</sup>

<sup>1</sup> Soil and Water Dept., Faculty of Agriculture, Kafrelsheikh University, Egypt (E-mail: hassan.elramady@agr.kfs.edu.eg)

<sup>2</sup> Plant Pathology Research Institute, Agriculture Research Center, Sakha, Kafr El-Sheikh, Egypt (E-mail: nagla\_abdelbaset@yahoo.com)

<sup>3</sup> Horticulture Dept., Faculty of Agriculture, Kafrelsheikh University, Egypt (E-mail: tashalaby@yahoo.com; ybayoumi2002@yahoo.com.sg)

<sup>4</sup> Agriculture Microbiology Department, Soil, Water and Environment Research Institute (SWERI), Sakha Agricultural Research Station, Agriculture Research Center (ARC), Kafr El-Sheikh 33717, Egypt; drelsakhawy@gmail.com; alaa.omara@yahoo.com)

<sup>5</sup> DE MÉK, Institute of Animal Science, Biotechnology and Nature Conservation, Department of Animal Husbandry, Nanofood Laboratory, Debrecen University, 4032 Debrecen, Hungary (E-mail: jprokisch@agr.unideb.hu)

<sup>6</sup> Physiology & Breeding of Horticultural Crops Laboratory, Horticulture Dept., Faculty of Agriculture, Kafrelsheikh University, Egypt

**T**HE MASSIVE application of nanoparticles in many sectors including medicine, agriculture and industry caused their inevitable release into different environmental compartments particularly groundwater and soil. The fate and behavior of these nanoparticles in soil and groundwater and their effects on soil biosystem remain largely unaddressed. Nano-selenium (Se-NPs) has intensive applications nowadays in different agroecosystems, but several studies, which highlight this behavior in plants and soils were carried out in individual case studies. In this concern, the interaction of nano-selenium with different nano-nutrients in soils and their uptake by cultivated plants still need urgent investigations. Based on the interaction is a common feature in the nature, there are still many unanswered questions regarding the interaction of Se-NPs with other nano-nutrients in soil for example: are these reactions synergistic or antagonistic? What are the factors controlling the bioavailability of nano-Se in soil in presence of other nano-nutrients? What is the expected role of stressful plants in orientation of this interaction among nano-nutrients?

**Keywords:** Biotic stress, Nano-silica, Plant pathogens, Nano-copper oxide, Nano-zinc oxide

#### Introduction

Plants can grow under different conditions including natural and stressful ones, but the productivity of cultivated plants definitely will be decreased due these stressors. Several studies have been published on the understanding response of

these stressful plants. These studies have increased our knowledge in understanding the plethora of biochemical, physiological, morphological, cellular and molecular responses under stressful plants (Saini et al. 2021). These stressors may include the abiotic stresses like salinity, drought, flooding, pollution, and heat stress; the biotic stress,

\*Corresponding author e-mail: ramady2000@gmail.com

Received 25/8/2021; Accepted 10/11/2021

DOI: 10.21608/jenvbs.2021.92788.1140

©2021 National Information and Documentation Center (NIDOC)

which result from the pathogens (Lassalle 2021). Under stressful conditions, many mechanisms or approaches take part to mitigate plant stress *via* endogenous agents such as phytohormones including cytokinin, gibberellin, ethylene, abscisic acid, brassinosteroids, salicylic acid and jasmonates (Saini *et al.* 2021) and/or exogenous factors like applied Se/nano-Se (Shalaby *et al.* 2021), Si/nano-Si (Akhtar *et al.* 2021), nano-TiO<sub>2</sub> (Ogunkunle *et al.* 2020; Shah *et al.* 2021), nano-ZnO (Ahmed *et al.* 2021a), nano-CuO (Joško *et al.* 2021), nano-iron/relatives (Zhang *et al.* 2021), and nano-organic fertilizer (Ahanger *et al.* 2021). As common studies, several investigations on individual nanoparticles on stressful plants have been published, but few studies were reported on combined nanoparticles (e.g., Ahmed *et al.* 2021a; Joško *et al.* 2021; Badawy *et al.* 2021).

Nano-selenium (i.e., nano-Se or Se-NPs) has unique properties like high bioavailability, low toxicity, high biological activity, high particle dispersion, and large surface area (Kumar and Prasad 2021). Thus, these nano-particles are widely in many sectors like medicine, e.g., drug delivery systems, medical diagnostics, antimicrobial agent, an antioxidant, anticancer agent, dietary supplement, and antidiabetic agent (Li and Xu 2020; He *et al.* 2021). In the agricultural sector, nano-Se has many beneficial effects on plant growth especially as nano-fertilizer (Gudkov *et al.* 2020), plant anti-stress (Shalaby *et al.* 2021), and nano-biofortification (El-Ramady *et al.* 2020, 2021a, b).

Therefore, this mini-review is an attempt to provide insight on the following themes (i) the distinguished role of nano-Se for decreasing plant stress, (ii) what is the fate and behavior of nano-nutrients in soils, and (iii) what is the expected interaction among nano-Se and other nano-nutrients? This is a call from the EBSS Editorial-board for more publications concerning this interaction among nano-nutrients in soil.

#### *Nano-Se for stressful plants*

A stress could be defined as “*the consequence of adverse effects of an external factor called stressor – or stress factor – on plant functions growth, and development*”. Cultivated crops face great challenges especially under environmental stresses, which decrease their productivity (Lassalle 2021). Environmental stressors include both abiotic stress (i.e., physical or chemical factors like salinity, drought, flooding, pollution) and biotic stress, which resulted from organisms

like plants, pathogens, insects (Othman *et al.* 2014; Lassalle 2021). Crop productivity has become limited under stressful conditions, which result from both the natural and anthropogenic origins (Lassalle *et al.* 2020). Many amendments have been used to mitigate biotic and abiotic stresses on cultivated plants particularly nanoparticles, e.g., Ag-NPs (Alabdallah and Hasan 2021), nano-TiO<sub>2</sub> (Ogunkunle *et al.* 2020) or nano-nutrients like nano-Se (Seliem *et al.* 2020; Qi *et al.* 2021), nano-silica (Lian *et al.* 2021), nano-CuO (Cota-Ruiz *et al.* 2020; Noman *et al.* 2020), nano-MgO (Ghassemi-Golezani *et al.* 2021), and nano-iron (Moradbeygi *et al.* 2020; Sreelakshmi *et al.* 2021).

Concerning nano-Se and its role to support the stressful plants, many studies confirmed this role, which is represented in the following investigations:

- 1- Enhancing peanuts plants (*Arachis hypogaea* L.) due to exogenous foliar application of nano-Se by activating antioxidant system under deficiency of nutrients in sandy soils (Hussein *et al.* 2019a).
- 2- Promoting the yield of pomegranate (*Punica granatum* L.) and its quality under arid zone (Zahedi *et al.* 2019b).
- 3- Alleviating Cd-toxicity in rice (*Oryza sativa* L.), Moldavian balm (*Dracocephalum moldavica* L.), and oilseed rape (*Brassica napus* L.) by modulating photosynthesis parameters and activities of antioxidant enzymes (Hussain *et al.* 2020; Azimi *et al.* 2021; Qi *et al.* 2021; Wang *et al.* 2021a).
- 4- Mitigating chilling stress on sugarcane (*Saccharum officinarum* L.) photosynthesis (Elsheery *et al.* 2020).
- 5- Protecting lettuce plants (*Lactuca sativa* L.) and strawberry (*Fragaria × ananassa* Duch) from salinity stress (Zahedi *et al.* 2019a; Soleymanzadeh *et al.* 2020).
- 6- Reducing the negative impacts of heat stress on *Chrysanthemum morifolium* Ramat by enhancing antioxidant enzyme activities particularly peroxidase and catalase at 150 mg nano-Se L<sup>-1</sup>; decreasing electrolyte leakage and polyphenol oxidase at 200 mg nano-Se L<sup>-1</sup> (Seliem *et al.* 2020).
- 7- Boosting the growth of cucumber (*Cucumis sativus* L.) and its productivity under soil salinity under heat stress by regulating the

osmotic balance and controlling stomatal opening through the high  $K^+$  content in cucumber leaves, which might support stressful cultivated plants (Shalaby et al. 2021).


- 8- Alleviating the salinity stress on bitter melon (*Momordica charantia*) plants by increasing proline,  $K^+$  and relative water content, the activity of antioxidant enzymes, and decreasing content of oxidants MDA and  $H_2O_2$  (Sheikhalipour et al. 2021).
- 9- Enhancing the quantity and quality of grains of wheat (*Triticum aestivum* L.) by 5–40%, under drought and heat stress as well as enhancing their tolerance to wheat crown and root rot diseases (El-Saadony et al. 2021a).

Many studies reported about the protective role of nanoparticles against many plant pathogens (i.e., bacteria, fungi, actinomycetes and nematode) like Ag-NPs (Gogoi et al. 2020; Javed et al. 2020; Paul and Roychoudhury 2021),  $SiO_2$ -NPs (Parveen and Siddiqui 2021), CuO-NPs (Ahmad et al. 2020; Mehta et al. 2020; Sathiyabama et al. 2020),  $La_2O_3$ -NPs (Adeel et al. 2021b), ZnO-NPs (Sharma et al. 2020; Khan and Siddiqui 2021), and carbon-based nanomaterials (Adeel et al. 2021a). On the other hand, few studies carried out on the role of nano-Se in supporting the cultivated plants under biotic stress like El-Saadony et al. (2021a), who investigated the wheat crown and root rot diseases induced by *Fusarium* spp. (i.e.,


*Fusarium culmorum*; *Fusarium graminearum*). Some studies also were reported on using of Se-NPs in controlling tomato leaf blight caused by *Alternaria alternata* under greenhouse conditions (El-Gazzar and Ismail 2020), and against tomato late blight disease (Joshi et al. 2021). More studies on stressful plants under biotic stress are needed including different phyto-diseases (e.g., leaf and root wilt, stem and leaf bright, leaf spot, stem canker, and soft rot). Some experiments were carried out in the Lab on nano-Se including *Alternaria solani* (Early blight) in tomato and potato (Fig. 1), *Rhizoctonia solani* (Fig. 2), and *Trichoderma* (Figs. 3 and 4).

#### Nano-nutrients in soils


Day by day, huge amounts of applied nanomaterials and/or nanoparticles are being used in different agroecosystems. The environmental content of nano-materials was predicted in soils to be within the ranged of  $10^{-7}$  to  $10^2$  mg  $kg^{-1}$ , and in landfills is thought to be about  $10^{-3}$  to  $10^3$  mg  $kg^{-1}$  (Adeel et al. 2021b). Several transformations could happen once nanoparticles release into soil, which may include adsorption, dissolution, sulphidation, aggregation and redox reactions (Joško et al. 2021). This release and dynamics of NPs in soil may depend mainly on characterization of NPs and soil, and may include soil pH, clay content, soil organic matter, cation exchange capacity and the rhizosphere characterization (Gao et al. 2017, 2019; Joško et al. 2020). The fate and behavior of nano-nutrients




**Fig. 1:** Effect of nano-selenium on *Alternaria solani* (Early blight) in tomato and potato was investigated on PDA media using 0, 25, and 50 mg  $l^{-1}$  nano-Se. These Petri dishes were infected by *Alternaria solani* and after 9 days from complete the growth in control petri dish, the reduction (inhibition rete) was calculated to be 25 and 75% for the 25 and 50 mg  $l^{-1}$  nano-Se, respectively



**Fig. 2:** Effect of nano-selenium on *Rhizoctonia solani* (as root rot or damping off or stem rot) was investigated on PDA media using 0, 50, 100 and 200 mg l<sup>-1</sup> nano-Se. These Petri dishes were infected by *Rhizoctonia solani* and after 4 days from complete the growth in control petri dish, the reduction (inhibition rete) was calculated to be 50, 50 and 75% for the 50, 100 and 200 mg l<sup>-1</sup> nano-Se, respectively



**Fig. 3:** Effect of nano-Se on *Trichoderma* spp. was very strange, where it was noticed that nano-Se changed the way of growing *Trichoderma* in the media and changed its behavior without inhibiting its growth. This may explain the reason of increasing the effectiveness of *Trichoderma*, but this needs further study for confirming



**Fig. 4:** Effect of nano-Se on the growth of *Trichoderma* spp. After 30 days from nano-Se treatment, *Trichoderma* changed the way of growing in the media, but changed its behavior without inhibiting its growth. This may help in increasing the effectiveness of *Trichoderma*.



or NPs in soil could be followed through three stations (*i*) nano-nutrients-soil interactions, (*ii*) nano-nutrients-rhizosphere interaction and (*iii*) nano-nutrients-plant interaction. Concerning the first station, soil environment can control the mobility of NPs, bioavailability and their toxicity in soil, which determine fate and sorption of nano-nutrients and subsequent impact on plants (Gao et al. 2018). Regarding nanoparticle-rhizosphere interaction, it is controlling by soil type (sandy, clayey, loamy) and time of exposure to NPs. In rhizosphere, different metabolites secreted (exudates) by plants and soil organisms and these exudates can modulate the fate of nano-nutrients (Gao et al. 2018). These root exudates are abundant mixtures in rhizosphere that control the behavior of NPs including organic acids, amino acids, sugars, and phenolics as low-molecular weight compounds and high-molecular weight compounds like mucilage (Gao et al. 2018; Wang et al. 2020). The third station (nano-nutrients-plant interaction), the phytotoxicity of nano-nutrients is governed by three factors including (1) plant type, species, and its growth stage, (2) nano-nutrients including size of NPs, concentration, and its aggregation and (3) growth media or experimental conditions including time, temperature, and method of exposure (Ahmed et al. 2021b).

Some distinguished attempts were already achieved on the fate and behavior of nano-nutrients or nanoparticles in soils and in some cases under cultivated plants, which could present in the following section:

- 1- The behavior of NPs and their bioavailability in soil under driven factors by rhizosphere processes, which include root exudates, the activities of soil microbes and earthworm, signal transduction as well as nano-nutrients absorption and their transport (Wang et al. 2020c).
- 2- Many kinds of nanoparticles have been discussed in the soil-plant environment including the interactions, assessing of their applications, fate and toxicity (Shrivastava et al. 2019; Rajput et al. 2020; Ameen et al. 2021; Rizwan et al. 2021) like iron oxide-NPs (Kamran et al. 2020), copper-based NPs (Bakshi and Kumar 2021), silver-NPs (Courtois et al. 2021), TiO<sub>2</sub>-NPs (Chavan et al. 2020), CeO<sub>2</sub>-NPs (Prakash et al. 2021), gold-NPs (Malejko et al. 2021), and nano-silica (Lian et al. 2021).

- 3- The expected interaction of nanoparticles with soil were reported by Ogunkunle et al. (2021). They discussed different NPs-sources and their transport into the soil with focus on NPs-characterization including their shape, surface chemistry, size, and water solubility, which may control the fate of NPs in soil and their various processes.
- 4- The release and behavior of nanoparticles in soils may follow one of the three following pathways for the interaction of NPs and soil components like (*i*) soil enzymes (de Oca-Vásquez et al. 2020; Mishra et al. 2021), (*ii*) soil properties particularly soil organic matter (Simonin et al. 2021), and (*iii*) soil macro-/microbial communities (Ameen et al. 2021; Macůrkova et al. 2021).

Concerning the efficacy of Se-NPs in soil, there are many multipurpose action and additional problems involved in the soil application of Se-NPs, which is related to content of soil organic matter (SOM) and its stimulation by the humic substance (HS) supplied simultaneously with Se and other nanoparticles. With respect that HS, Se-NPs, and SOM functioning in the soil are linked to the degradation of organic matter, which is mainly controlled by soil microbial activities and linked to the soil fertility (Gudkov et al. 2020). The proper Se-NPs functioning in soil is controlling by soil tillage, ploughing and other soil cultivation procedures, which may be linked to formation of soil aggregates system. Soil moisture content also is an important factor controlling the functions of Se-NPs, HS, poly-microbial biofilms and the plant growth. The high soil water content (flooding) may cause a problem in leaching the nanoparticles and other soil matters (Gudkov et al. 2020). The profile distribution of natural NPs in paddy soils may affect by long-term rice cultivation (Huang et al. 2021).

Some published studies focused on responses of different cultivated crops to many different nanoparticles like applied NPs of Al<sub>2</sub>O<sub>3</sub>, CuO, TiO<sub>2</sub>, ZnO to cucumber plants cultivated on sandy-clay-loam field soil (Ahmed et al. 2021c). Many metal-oxide NPs are transformed by environmental factors such as CuO-NPs, which could be degraded into copper sulfate hydroxides and then release Cu<sup>2+</sup> that increases the bioavailability of Cu in soil and its bio-uptake by cultivated plants (Servin et al. 2017). Concerning the uptake of nano-Se, cultivated plants can take up the Se-NPs, which can influx

into roots as a “*passive diffusion process*” and the uptake rate of nano-Se (chemical and biological forms) is lower than for the selenite form (Hu *et al.* 2018). The chemical nano-Se could easily uptake by plant roots more than the biological one especially when the nanoparticles are below 50 nm in size. Few nanoparticles indicate that Se could be transported from plant roots into shoots, and then they will rapidly assimilate into selenite and organic forms in roots and shoots. The uptake of Bio-Se-NPs could be inhibited by aquaporins inhibitor in a higher rate compared to chemical Se-NPs in 93.4 and 92.5%, respectively (Hu *et al.* 2018). In general, the uptake rate of selenite and selenate was approximately equal, and both exhibited higher uptake rate than nano-Se, which was estimated, in case of selenite, by 2.5-fold higher than nano-Se (Hu *et al.* 2018). Concerning the transformation of nano-Se in roots, it is found that selenite was rapidly assimilated in wheat roots into organic forms (e.g., SeMet), whereas the nano-Se appeared stable in the solution and did not oxidize to selenite (Hu *et al.* 2018). The uptake of Se-NPs rice roots was confirmed; then they can transport successfully into the aerial parts and the dominant Se-species was SeMet under different Se-NPs treatment (Wang *et al.* 2020b). Selenium is characterized by its ability to be lost via volatilization into the atmosphere such as H<sub>2</sub>Se, DMSe, DMDSe, DMS<sub>2</sub>S, and DMS<sub>2</sub>O<sub>2</sub>, where Se-species “DMSe” is the main dominant volatile Se species (Ye *et al.* 2021). The great role of microorganisms in Se is to biosynthesize the nano-Se (11 – 700 nm in diameter), whereas the biosynthesized organic form by fungi ranges from 17 to 150 nm (Wang *et al.* 2022). Red elemental nano-Se spheres in water may produce H<sub>2</sub>Se and H<sub>2</sub>SeO<sub>3</sub> in small amount according to the following equation  $3\text{Se} + 3\text{H}_2\text{O} \leftrightarrow 2\text{H}_2\text{Se} + \text{H}_2\text{SeO}_3$ . Afterwards, H<sub>2</sub>Se and H<sub>2</sub>SeO<sub>3</sub> may react together and precipitate forming crystals when the solution (El-Ramady *et al.* 2015).

#### *Nano-Se and its interaction with other nano-nutrients*

Several studies have been published about the individual impacts of nanoparticles of different nutrients on cultivated crops like nano-Se (Shalaby *et al.* 2021), nano-silica (Mathur and Roy 2020; Lian *et al.* 2021), nano-Cuo (Li *et al.* 2021), nano-Fe<sub>2</sub>O<sub>3</sub> (Zhang *et al.* 2021), and nano-ZnO (Keerthana *et al.* 2021), but few studies focused on the combined impact of two or more nano-nutrients (**Table 1**). The co-existence of different engineered nano-nutrients in the agro-

environment is an emerging issue remaining poorly investigated (Joško *et al.* 2021). **Table 2** presents some published studies to be compared nano-Se and its interaction with some other nano-nutrients.

Many of these nano-nutrients applied to mitigate the plant stressful conditions like drought (Zahedi *et al.* 2020), heavy metals stress (Hussain *et al.* 2020; Wang *et al.* 2020a), and soil salinity (Badawy *et al.* 2021). Concerning the heavy metals pollution, foliar application of the combined nano-Se and nano-Si has shown success in decreasing the accumulation of Cd and Pb metals in rice grains, hence improved the rice grain quality and Se-biofortification (Hussain *et al.* 2020). Some mixtures of nano-nutrients like nano-ZnO and nano-CuO caused a reduction in the inhibition of root elongation of cress, cucumber, flax, and wheat comparing with single exposure to these particles (Joško *et al.* 2017). On the other hand, some studies reported the opposite trend like a higher reduction in spinach biomass irrigated with mixtures of nano-CuO and nano-ZnO than individual treatments (Singh and Kumar 2020). The bioactivity of these nano-nutrients may be attributed to nature of these nano-nutrients (García-Gomez *et al.* 2017). In other study, the mixtures of nano-CuO and nano-La<sub>2</sub>O<sub>3</sub> decreased the accumulation of Cu in zucchini plant tissues than single exposure to its NPs (Pagano *et al.* 2017). So, several studies demonstrated the changes in the nutritional profile of many plants exposed to different-NPs (Cota-Ruiz *et al.* 2020; Sharifan *et al.* 2020). Therefore, an urgent question arises then about how the mineral composition of plants will be affected by co-existing these nano-nutrients (Joško *et al.* 2021).

Combined application of nano-Se and nano-SiO<sub>2</sub> at both 50 and 100 mg L<sup>-1</sup> improved the yield and fruit quality of strawberry plants compared to the control which was grown under normal and drought stress conditions, which included normal irrigation at 100% Field Capacity (FC), moderate stress at 60% FC, and severe stress at 25% FC. The highest dose of applied nano-Se and nano-SiO<sub>2</sub> (100 mg L<sup>-1</sup>) preserved more of their photosynthetic pigments compared with other treated plants and presented higher levels of key osmolytes such as carbohydrate and proline. Applied nano-Se/SiO<sub>2</sub> improved strawberry fruit quality and nutritional value (total phenolic compounds, anthocyanin, vitamin C and antioxidant activity) under drought stress (Zahedi

**TABLE 1: The most important nano-nutrients already used in crop production and some details concerning cultivated crop, and some individual and combined published studies**

Nano-nutrient alone or and its coexistence nano-nutrient	Cultivated crop and its scientific name	References	
<i>Some published studies on individual nano-nutrients</i>			
<b>Nano-Se</b>	Cucumber ( <i>Cucumis sativus</i> L.)	Shalaby et al. (2021)	
	Chicory ( <i>Cichorium intybus</i> L.)	Abedi et al. (2021)	
	Wheat ( <i>Triticum aestivum</i> L.)	El-Saadony et al. (2021a)	
	Lemon balm ( <i>Melissa officinalis</i> L.)	Ghasemian et al. (2021)	
	Common bean ( <i>Phaseolus vulgaris</i> )	Rady et al. (2021)	
	Rice ( <i>Oryza sativa</i> L.)	Wang et al. (2020b)	
	Groundnut ( <i>Arachis hypogaea</i> L.)	Hussein et al. (2019a, b)	
	Tomato ( <i>Solanum lycopersicum</i> L.)	Neysanian et al. (2020)	
<b>Nano-Cu or CuO</b>	Tomato ( <i>Solanum lycopersicum</i> L.)	Morales-Espinoza et al. (2019)	
	Rape ( <i>Brassica napus</i> L.)	Li et al. (2021)	
	Alfalfa ( <i>Medicago sativa</i> L.)	Cota-Ruiz et al. (2020)	
	Maize ( <i>Zea mays</i> L.)	Pu et al. (2019)	
<b>Nano-Fe and relatives</b>	Bell pepper ( <i>Capsicum annuum</i> )	Rawat et al. (2019)	
	Rice ( <i>Oryza sativa</i> L.)	Zhang et al. (2021)	
	Fever nut ( <i>Caesalpinia bonducella</i> )	Khalid et al. (2021)	
	Foxtail millet ( <i>Setaria italica</i> )	Sreelakshmi et al. (2021)	
<b>Nano-MgO</b>	nZVI Rice ( <i>Oryza sativa</i> L.)	Guha et al. (2020)	
	Sweet basil ( <i>Ocimum basilicum</i> L.)	Tavallali et al. (2020)	
	Fever nut ( <i>Caesalpinia bonducella</i> )	Khalid et al. (2021)	
<b>Nano-Si or silica</b>	Cowpea ( <i>Vigna unguiculata</i> L.)	Tauseef et al. (2021)	
	Wheat ( <i>Triticum aestivum</i> L.)	Akhtar et al. (2021).	
	Rice ( <i>Oryza sativa</i> L.)	Banerjee et al. (2021)	
	Common bean ( <i>Phaseolus vulgaris</i> )	El-Saadony et al. (2021b)	
<b>Nano-Zn or ZnO</b>	Lemon balm ( <i>Melissa officinalis</i> L.)	Hatami et al. (2021)	
	Maize ( <i>Zea mays</i> L.)	Kumaraswamy et al. (2021)	
	Lettuce ( <i>Lactuca sativa</i> L.)	Lian et al. (2021)	
	Okra ( <i>Abelmoschus esculentus</i> L.)	Keerthana et al. (2021)	
	Wheat ( <i>Triticum aestivum</i> L.)	Rai-Kalal et al. (2021)	
<i>Some published studies on combined nano-nutrients</i>	Onion ( <i>Allium cepa</i> L.)	Debnath et al. (2020)	
	Rice ( <i>Oryza sativa</i> L.)	Chutipaijit et al. (2018)	
	Nano-CuO and nano-ZnO	Maize ( <i>Zea mays</i> L.)	Ahmed et al. (2021a)
	nano-CuO and nano-ZnO	Barley ( <i>Hordeum vulgare</i> L.)	Joško et al. (2021)
	Nano-Se and nano-ZnO	Lemon balm ( <i>Melissa officinalis</i> L.)	Babajani et al. (2019)
	Nano-Se and Nano-Si	Rice ( <i>Oryza sativa</i> L.)	Badawy et al. (2021)
	Nano-Se and nano-copper	Tomato ( <i>Solanum lycopersicum</i> L.)	Hernández-Hernández et al. (2019)
	Nano-Se and nano-copper	Tomato ( <i>Solanum lycopersicum</i> L.)	Quiterio-Gutiérrez et al. (2019)
	Nano-Se and Nano-Si	Rice ( <i>Oryza sativa</i> L.)	Hussain et al. (2020)
	Nano-Se and Nano-silica	Strawberry ( <i>Fragaria × ananassa</i> Duch.)	Zahedi et al. (2020)

Note: nZVI: nano-scale zero valent iron

**TABLE 2: Some published case studies of the interaction of nano-Se and other nano-nutrients applied for cultivated plants**

Cultivated plant and its scientific name	Nano-Se and its applied dose	Applied nano-nutrient and its details	Purpose of the study	Reference
Rice ( <i>Oryza sativa</i> L.)	Chemical nano-Se (12.26 nm) at 5, 10 and 20 mg L <sup>-1</sup>	Nano-silica (18.04 nm) at 5, 10 and 20 mg L <sup>-1</sup>	Foliar applied of both NPs regarded the accumulation of Cd and Pb in rice grains	Hussain et al. (2020)
Rice ( <i>Oryza sativa</i> L.)	Se-NPs at 4, 6, and 12 mg L <sup>-1</sup> (75 nm)	Si-NPs at 15, 22, and 44 mg L <sup>-1</sup> (6 and 7 nm)	Foliar applied Si and Se-NPs reduced Cd and Pb translocation to rice grains; mitigate oxidative damage	Wang et al. (2020a)
Strawberry ( <i>Fragaria × ananassa</i> Duch.)	Chemical Se-NPs (10-45 nm) in combined with nano-silica	Spraying SiO <sub>2</sub> -NPs (20-30 nm), SiO <sub>2</sub> , Se, and nano-Se/SiO <sub>2</sub> at 50, 100 mg l <sup>-1</sup>	Both nano-Se and nano-silica supported cultivated strawberry under drought stress	Zahedi et al. (2020)
Lemon balm ( <i>Melissa officinalis</i> )	Chemical nano-Se at 10, and 50 mg l <sup>-1</sup> (10-45 nm)	Nano-ZnO (10–30 nm) at 100, and 300 mg l <sup>-1</sup>	Possible impacts of both nano-nutrients on growth, toxicity, and antioxidants in plants	Babajani et al. (2019)
Tomato ( <i>Solanum lycopersicum</i> L.)	Chemical Se-NPs at 1, 10, 20 mg L <sup>-1</sup> (2-20 nm)	Cu NPs at 10, 50, 250 mg L <sup>-1</sup> (42 nm)	Combined nano-nutrients increased yield and fruit quality of tomato	Hernández-Hernández et al. (2019)
Tomato ( <i>Solanum lycopersicum</i> L.)	Chemical Se-NPs at 10 and 20 mg L <sup>-1</sup> (2–20 nm)	Cu-NPs at 10 and 50 mg L <sup>-1</sup> (50 nm)	Applied nanoparticles increasing antioxidant capacity under early blight disease stress	Quiterio-Gutiérrez et al. (2019)

et al. 2020).

On the other hand, the diverse aspects of the interplay between different-NPs and cultivated plants are very important for discussing this interplay (Hu and Xianyu 2021). Therefore, several questions could be asked concerning the combined application of nano-nutrients and the expected interaction like: what will be happened when the nano-nutrients apply as binary mixtures? Is this mixture of anno-nutrients synergistic or antagonistic for plant growth? To what extent we can expect the interactions of combined exposure of nano-nutrients, which may include adsorption and/or competition between ionic and particulate metals as well as with biomolecules?

### Conclusions

The rising wide-scale uses of Se-NPs for different applications in agriculture, industry, and environmental issues would undoubtedly lead to their dissemination in the agroecosystems (mainly groundwater and soil). Based on the

environmental impacts of Se-NPs are speedily progressing in individual cases, some major understanding is needed on Se-NPs mechanism of their interactions with other nano-nutrients in different soil components and their subsequent impacts on plants and microorganisms. Several nanoparticles already were investigated for their fate, behavior and impacts on the soil-plant system, but the combined and multi-impacts nano-nutrients still need further investigations. These multi-interaction cases in soil might influence by the functioning of the soil nutrient cycles, soil microbial activity, different reactions in the rhizosphere, plant exudates, and soil properties. Concerning the bio-transformation of Se-NPs or their chemical complexes in soil, this important area still needs future investigations. The phytotoxicity of Se-NPs and its uptake might vary in different plant species and the structure of exposure media. Several open questions are still needed to be investigated such as what is the expected mechanism of nano-nutrients and



their role in rhizosphere and their interactions in influencing Se-NPs plant uptake and translocation? What are the expected and different pathways of nano-Se interactions with other nano-nutrients in soil based on their individual properties?

*Ethics approval and consent to participate:*

This article does not contain any studies with human participants or animals performed by any of the authors.

*Consent for publication:*

All authors declare their consent for publication.

*Funding:* This research received no external funding.

*Author contribution:*

This project was conceived and led by El-Ramady. All authors have read and agreed to the published version of the manuscript.

*Conflicts of Interest:*

The author declares no conflict of interest.

**Acknowledgments:**

This work was financialized and supported by the Central Department of Mission, Egyptian Ministry of Higher Education (Mission 19/2020) for H. El-Ramady and also by the Hungarian Tempus Public Foundation (TPF), grant no. AK-00152-002/2021.

**References**

- Abedi S, Iranbakhsh A, Ardebili ZO, Ebadi M (2021). Nitric oxide and selenium nanoparticles confer changes in growth, metabolism, antioxidant machinery, gene expression, and flowering in chicory (*Cichorium intybus* L.): potential benefits and risk assessment. *Environmental Science and Pollution Research* 28, 3136–3148. <https://doi.org/10.1007/s11356-020-10706-2>
- Adeel M, Farooq T, White JC, Hao Y, He Z, Rui Y (2021a). Carbon-based nanomaterials suppress Tobacco Mosaic Virus (TMV) infection and induce resistance in *Nicotiana benthamiana*. *J. Hazard. Mater.* 404, 124167. <https://doi.org/10.1016/j.jhazmat.2020.124167>
- Adeel M, Shakoor N, Hussain T, Azeem I, Zhou P, Zhang P, Hao Y, Rinklebe J, Rui Y (2021b). Bio-interaction of nano and bulk lanthanum and ytterbium oxides in soil system: Biochemical, genetic, and histopathological effects on *Eisenia fetida*. *Journal of Hazardous Materials* 415, 125574. <https://doi.org/10.1016/j.jhazmat.2021.125574>
- Ahanger MA, Qi M, Huang Z, Xu X, Begum N, Qin C, Zhang C, Ahmad N, Mustafa NS, Ashraf M, Zhang L (2021). Improving growth and photosynthetic performance of drought stressed tomato by application of nano-organic fertilizer involves up-regulation of nitrogen, antioxidant and osmolyte metabolism. *Ecotoxicology and Environmental Safety* 216, 112195. <https://doi.org/10.1016/j.ecoenv.2021.112195>
- Ahmad H, Venugopal K, Bhat AH, Kavitha K, Ramanan A, Rajagopal K, Srinivasan R, Manikandan E (2020). Enhanced Biosynthesis Synthesis of Copper Oxide Nanoparticles (CuO-NPs) for their Antifungal Activity Toxicity against Major Phyto-Pathogens of Apple Orchards. *Pharm Res*, 37, 246. <https://doi.org/10.1007/s11095-020-02966-x>
- Ahmed B, Rizvi A, Ali K, Lee J, Zaidi A, Khan MS, Musarrat J (2021b). Nanoparticles in the soil–plant system: a review. *Environmental Chemistry Letters* 19, 1545–1609. <https://doi.org/10.1007/s10311-020-01138-y>
- Ahmed B, Rizvi A, Syed A, Elgorban AM, Khan MS, AL-Shwaiman HA, Musarrat J, Lee J (2021a). Differential responses of maize (*Zea mays*) at the physiological, biomolecular, and nutrient levels when cultivated in the presence of nano or bulk ZnO or CuO or Zn<sup>2+</sup> or Cu<sup>2+</sup> ions. *Journal of Hazardous Materials* 419, 126493. <https://doi.org/10.1016/j.jhazmat.2021.126493>
- Ahmed B, Rizvi A, Syed A, Jailani A, Elgorban AM, Khan MS, AL-Shwaiman HA, Lee J (2021c). Differential bioaccumulations and ecotoxicological impacts of metal-oxide nanoparticles, bulk materials, and metal-ions in cucumbers grown in sandy clay loam soil. *Environmental Pollution* 289, 117854. <https://doi.org/10.1016/j.envpol.2021.117854>
- Akhtar N, Ilyas N, Mashwani Z, Hayat R, Yasmin H, Noureldeen A, Ahmad P (2021). Synergistic effects of plant growth promoting rhizobacteria and silicon dioxide nano-particles for amelioration of drought stress in wheat. *Plant Physiology and Biochemistry* 166, 160–176. <https://doi.org/10.1016/j.plaphy.2021.05.039>
- Alabdallah NM, Hasan MM (2021). Plant-based green synthesis of silver nanoparticles and its effective role in abiotic stress tolerance in crop plants. *Saudi Journal of Biological Sciences*, <https://doi.org/10.1016/j.sjbs.2021.05.081>
- Ameen F, Alsamhary K, Alabdullatif JA, ALNadhari

- S (2021). A review on metal-based nanoparticles and their toxicity to beneficial soil bacteria and fungi. *Ecotoxicology and Environmental Safety* 213, 112027. <https://doi.org/10.1016/j.ecoenv.2021.112027>
- Azimi F, Oraei M, Gohari G, Panahirad S, Farmarzi A (2021). Chitosan-selenium nanoparticles (Cs–Se NPs) modulate the photosynthesis parameters, antioxidant enzymes activities and essential oils in *Dracocephalum moldavica* L. under cadmium toxicity stress. *Plant Physiology and Biochemistry* 167, 257–268. <https://doi.org/10.1016/j.plaphy.2021.08.013>
- Babajani A, Iranbakhsh A, Ardebili ZO, Eslami B (2019). Differential growth, nutrition, physiology, and gene expression in *Melissa officinalis* mediated by zinc oxide and elemental selenium nanoparticles. *Environmental Science and Pollution Research* 26, 24430–24444. <https://doi.org/10.1007/s11356-019-05676-z>
- Badawy SA, Zayed BA, Bassiouni SMA, Mahdi AHA, Majrashi A, Ali EF, Seleiman MF (2021). Influence of Nano Silicon and Nano Selenium on Root Characters, Growth, Ion Selectivity, Yield, and Yield Components of Rice (*Oryza sativa* L.) under Salinity Conditions. *Plants*, 10, 1657. <https://doi.org/10.3390/plants10081657>
- Bakshi M, Kumar A (2021). Copper-based nanoparticles in the soil-plant environment: Assessing their applications, interactions, fate and toxicity. *Chemosphere* 281, 130940. <https://doi.org/10.1016/j.chemosphere.2021.130940>
- Banerjee A, Singh A, Sudarshan M, Roychoudhury A (2021). Silicon nanoparticle-pulsing mitigates fluoride stress in rice by finetuning the ionic and metabolomic balance and refining agronomic traits. *Chemosphere* 262, 127826. <https://doi.org/10.1016/j.chemosphere.2020.127826>
- Chavan S, Sarangdhar V, Nandanathangam V (2020). Toxicological effects of TiO<sub>2</sub> nanoparticles on plant growth promoting soil bacteria. *Emerging Contaminants* 6, 87-92. <https://doi.org/10.1016/j.emcon.2020.01.003>
- Chutipaijit S, Whalley AJS, Sutjaritvorakul T (2018). In vitro plant growth promotion by ZnO nanomaterials in indica rice seedlings (*Oryza sativa* L.). *Materials Today: Proceedings* 5, 14944–14949.
- Cota-Ruiz K, Ye Y, Valdes C, Deng C, Wang Y, Hernández-Viezcás JA, Duarte-Gardea M, Gardea-Torresdey JL (2020). Copper nanowires as nanofertilizers for alfalfa plants: Understanding nano-bio systems interactions from microbial genomics, plant molecular responses and spectroscopic studies. *Science of the Total Environment* 742, 140572. <https://doi.org/10.1016/j.scitotenv.2020.140572>
- Courtois P, de Vaufléury A, Grosser A, Lors C, Vandenbulcke F (2021). Transfer of sulfidized silver from silver nanoparticles, in sewage sludge, to plants and primary consumers in agricultural soil environment. *Science of the Total Environment* 777, 145900. <https://doi.org/10.1016/j.scitotenv.2021.145900>
- de Oca-Vásquez GM, Solano-Campos F, Vega-Baudrit JR, López-Mondéjar R, Odriozola I, Vera A, Moreno JL, Bastida F (2020). Environmentally relevant concentrations of silver nanoparticles diminish soil microbial biomass but do not alter enzyme activities or microbial diversity. *Journal of Hazardous Materials* 391, 122224. <https://doi.org/10.1016/j.jhazmat.2020.122224>
- Debnath P, Mondal A, Sen K, Mishra D, Mondal NK (2020). Genotoxicity study of nano Al<sub>2</sub>O<sub>3</sub>, TiO<sub>2</sub> and ZnO along with UV-B exposure: An *Allium cepa* root tip assay. *Science of the Total Environment* 713, 136592. <https://doi.org/10.1016/j.scitotenv.2020.136592>
- El-Gazzar N, Ismail AM (2020). The potential use of Titanium, Silver and Selenium nanoparticles in controlling leaf blight of tomato caused by *Alternaria alternata*. *Biocatalysis and Agricultural Biotechnology* 27, 101708. <https://doi.org/10.1016/j.bcab.2020.101708>
- El-Ramady H, Abdalla N, Elbasiouny H, Elbehiry F, Elsakhawy T, Omara AED, Amer M, Bayoumi Y, Shalaby TA, Eid Y, Zia-ur-Rehman M (2021a). Nano-biofortification of different crops to immune against COVID-19: A review. *Ecotoxicology and Environmental Safety*, 222, 112500. <https://doi.org/10.1016/j.ecoenv.2021.112500>
- El-Ramady H, El-Mahdy S, Awad A, Nassar S, Osman O, Metwally E, Aly E, Fares E, El-Henawy A (2021b). Is nano-biofortification the right approach for malnutrition in the era of COVID-19 and climate change? *Egypt. J. Soil Sci.*, 61 (2), 141-150, DOI 10.21608/ejss.2021.75653.1445
- El-Ramady H, Faizy SE-D, Abdalla N, Taha H, Domokos-Szabolcsy É, Fari M, Elsakhawy T, Omara AE-D, Shalaby T, Bayoumi Y, Shehata S, Geilfus C-M, Brevik EC (2020). Selenium and

- Nano-Selenium Biofortification for Human Health: Opportunities and Challenges. *Soil Syst.*, 4, 57. doi:10.3390/soilsystems4030057
- El-Ramady HR, Domokos-Szabolcsy É, Shalaby TA, Prokisch J, Fári M (2015). Selenium in Agriculture: Water, Air, Soil, Plants, Food, Animals and Nanoselenium. In: E. Lichtfouse et al. (eds.), CO<sub>2</sub> Sequestration, Biofuels and Depollution, Environmental Chemistry for a Sustainable World 5, DOI 10.1007/978-3-319-11906-9\_5, pp: 153 – 232. Springer International Publishing Switzerland
- El-Saadony MT, Desoky EM, Saad AM, Eid RSM, Selem E, Elrys AS (2021b). Biological silicon nanoparticles improve *Phaseolus vulgaris* L. yield and minimize its contaminant contents on a heavy metals-contaminated saline soil. *Journal of Environmental Sciences* 106, 1–14. <https://doi.org/10.1016/j.jes.2021.01.012>
- El-Saadony MT, Saad MM, Najjar AA, Alzahrani SO, Alkhatib FM, Shafi ME, Selem E, Desoky EM, Fouda SEE, El-Tahan AM, Hassan MAA (2021a). The use of biological selenium nanoparticles to suppress *Triticum aestivum* L. crown and root rot diseases induced by Fusarium species and improve yield under drought and heat stress. *Saudi Journal of Biological Sciences* 28, 4461–4471. <https://doi.org/10.1016/j.sjbs.2021.04.043>
- Gao X, Avellan A, Laughton S, Vaidya R, Rodrigues SM, Casman EA, Lowry GV (2018). CuO nanoparticle dissolution and toxicity to wheat (*Triticum aestivum*) in rhizosphere soil. *Environ. Sci. Technol.* 52, 2888–2897. <https://doi.org/10.1021/acs.est.7b05816>
- Gao X, Rodrigues SM, Spielman-Sun E, Lopes S, Rodrigues S, Zhang Y, Avellan A, Duarte RMBO, Duarte A, Casman EA, Lowry GV (2019). Effect of soil organic matter, soil pH, and moisture content on solubility and dissolution rate of CuO NPs in soil. *Environ. Sci. Technol.* 53, 4959–4967. <https://doi.org/10.1021/acs.est.8b07243>
- Gao X, Spielman-Sun E, Rodrigues SM, Casman EA, Lowry GV (2017). Time and nanoparticle concentration affect the extractability of Cu from CuO NP-amended soil. *Environ. Sci. Technol.* 51, 2226–2234. <https://doi.org/10.1021/acs.est.6b04705>
- García-Gomez C, Obrador A, Gonzalez D, Babin M, Fernandez MD (2017). Comparative effect of ZnO NPs, ZnO bulk and ZnSO<sub>4</sub> in the antioxidant defenses of two plant species growing in two agricultural soils under greenhouse conditions. *Sci. Total Environ.* 589, 11–24. <https://doi.org/10.1016/j.scitotenv.2017.02.153>
- Ghasemian S, Masoudian N, Nematpour FS, Afshar AS (2021). Selenium nanoparticles stimulate growth, physiology, and gene expression to alleviate salt stress in *Melissa officinalis*. *Biologia*, <https://doi.org/10.1007/s11756-021-00854-2>
- Ghassemi-Golezani K, Farhangi-Abriz S (2021). Biochar-based metal oxide nanocomposites of magnesium and manganese improved root development and productivity of safflower (*Carthamus tinctorius* L.) under salt stress. *Rhizosphere* 19, 100416. <https://doi.org/10.1016/j.rhisph.2021.100416>
- Gogoi B, Kumar R, Upadhyay J, Borah D (2020). Facile biogenic synthesis of silver nanoparticles (AgNPs) by *Citrus grandis* (L.) Osbeck fruit extract with excellent antimicrobial potential against plant pathogens. *SN Applied Sciences* 2, 1723. <https://doi.org/10.1007/s42452-020-03529-w>
- Gudkov SV, Shafeev GA, Glinushkin AP, Shkirin AV, Barmina EV, Rakov II, Simakin AV, Kislov AV, Astashev ME, Vodeneev VA, Kalinitchenko VP (2020). Production and Use of Selenium Nanoparticles as Fertilizers. *ACS Omega.* 5(28), 17767–17774. Doi: 10.1021/acsomega.0c02448
- Guha T, Gopal G, Chatterjee R, Mukherjee A, Kundu R (2020). Differential growth and metabolic responses induced by nano-scale zero valent iron in germinating seeds and seedlings of *Oryza sativa* L. cv. Swarna. *Ecotoxicology and Environmental Safety* 204, 111104. <https://doi.org/10.1016/j.ecoenv.2020.111104>
- Hatami M, Khanizadeh P, Bovand F, Aghae A (2021). Silicon nanoparticle-mediated seed priming and Pseudomonas spp. inoculation augment growth, physiology and antioxidant metabolic status in *Melissa officinalis* L. plants. *Industrial Crops & Products* 162, 113238. <https://doi.org/10.1016/j.indcrop.2021.113238>
- He L, Zhao J, Wang L, Liu Q, Fan Y, Li B, Yu Y-L, Chen C, Lia Y-F (2021). Using nano-selenium to combat Coronavirus Disease 2019 (COVID-19)? *Nano Today* 36, 101037. <https://doi.org/10.1016/j.nantod.2020.101037>
- Hernández-Hernández H, Quiterio-Gutiérrez T, Cadenas-Pliego G, Ortega-Ortiz H, Hernández-Fuentes AD, Cabrera de la Fuente M, Valdés-Reyna J, Juárez-Maldonado A (2019). Impact of selenium

- and copper nanoparticles on yield, antioxidant system, and fruit quality of tomato plants. *Plants* 8, 355. DOI: 10.3390/plants8100355
- Hu J, Xianyu Y (2021). When nano meets plants: A review on the interplay between nanoparticles and plants. *Nano Today* 38, 101143. <https://doi.org/10.1016/j.nantod.2021.101143>
- Hu T, Li H, Li J, Zhao G, Wu W, Liu L, Wang Q, Guo Y (2018). Absorption and Bio-Transformation of Selenium Nanoparticles by Wheat Seedlings (*Triticum aestivum* L.). *Front. Plant Sci.* 9:597. doi: 10.3389/fpls.2018.00597
- Huang D, Zhu X, Xu B, He Y, Zhang M, Liu F, Lian Z, Dahlggen RA, PC Brookes PC, Xu J (2021). Changes in profile distribution and chemical properties of natural nanoparticles in paddy soils as affected by long-term rice cultivation. *Pedosphere* 31(5), 659–669. doi:10.1016/S1002-0160(21)60015-2
- Hussain B, Lin Q, Hamid Y, Sanaullah M, Di L, Hashmi ML, Khan MB, He Z, Yang X (2020). Foliage application of selenium and silicon nanoparticles alleviates Cd and Pb toxicity in rice (*Oryza sativa* L.). *Science of the Total Environment* 712, 136497. <https://doi.org/10.1016/j.scitotenv.2020.136497>
- Hussein H-AA, Darwesh OM, Mekki BB (2019b). Environmentally friendly nano-selenium to improve antioxidant system and growth of groundnut cultivars under sandy soil conditions. *Biocatalysis and Agricultural Biotechnology* 18, 101080. <https://doi.org/10.1016/j.bcab.2019.101080>
- Hussein H-AA, Darwesh OM, Mekki BB, El-Hallouty SM (2019a). Evaluation of cytotoxicity, biochemical profile and yield components of groundnut plants treated with nano-selenium. *Biotechnology Reports* 24, e00377. <https://doi.org/10.1016/j.btre.2019.e00377>
- Javed B, Raja NI, Nadhman A, Mashwani Z (2020). Understanding the potential of biofabricated nonoxidative silver nanoparticles to eradicate Leishmania and plant bacterial pathogens. *Applied Nanoscience* 10, 2057–2067. <https://doi.org/10.1007/s13204-020-01355-5>
- Joshi SM, De Britto S, Jogaiah S (2021). Myco-engineered selenium nanoparticles elicit resistance against tomato late blight disease by regulating differential expression of cellular, biochemical and defense responsive genes. *Journal of Biotechnology* 325, 196–206. <https://doi.org/10.1016/j.jbiotec.2020.10.023>
- Joško I, Dobrzynska J, Dobrowolski R, Kusiak M, Terpiłowski K (2020). The effect of pH and ageing on the fate of CuO and ZnO nanoparticles in soils. *Sci. Total Environ.* 721, 137771 <https://doi.org/10.1016/j.scitotenv.2020.137771>
- Joško I, Kusiak M, Xing B, Oleszczuk P (2021). Combined effect of nano-CuO and nano-ZnO in plant-related system: From bioavailability in soil to transcriptional regulation of metal homeostasis in barley. *Journal of Hazardous Materials* 416, 126230. <https://doi.org/10.1016/j.jhazmat.2021.126230>
- Joško I, Oleszczuk P, Skwarek E (2017). Toxicity of combined mixtures of nanoparticles to plants. *J. Hazard. Mater.* 331, 200–209. <https://doi.org/10.1016/j.jhazmat.2017.02.028>
- Kamran M, Ali H, Saeed MF, Bakhat HF, Hassan Z, Tahir M, Abbas G, Naeem MA, Rashid MI, Shah GM (2020). Unraveling the toxic effects of iron oxide nanoparticles on nitrogen cycling through manure-soil-plant continuum. *Ecotoxicology and Environmental Safety* 205, 111099. <https://doi.org/10.1016/j.ecoenv.2020.111099>
- Keerthana P, Vijayakumar S, Vidhya E, Punitha VN, Nilavukkarasi M, Praseetha PK (2021). Biogenesis of ZnO nanoparticles for revolutionizing agriculture: A step towards anti-infection and growth promotion in plants. *Industrial Crops & Products* 170, 113762. <https://doi.org/10.1016/j.indcrop.2021.113762>
- Khalid U, Sher F, Noreen S, Lima EC, Rasheed T, Sehar S, Amami R (2021). Comparative effects of conventional and nano-enabled fertilizers on morphological and physiological attributes of *Caesalpinia bonducella* plants, *Journal of the Saudi Society of Agricultural Sciences*, <https://doi.org/10.1016/j.jssas.2021.06.011>
- Khan MR, Siddiqui ZA (2021). Role of zinc oxide nanoparticles in the management of disease complex of beetroot (*Beta vulgaris* L.) caused by *Pectobacterium betavasculorum*, *Meloidogyne incognita* and *Rhizoctonia solani*. *Horticulture, Environment, and Biotechnology* 62, 225–241. <https://doi.org/10.1007/s13580-020-00312-z>
- Kumar A, KS Prasad (2021). Role of nano-selenium in health and environment. *Journal of Biotechnology* 325, 152–163. <https://doi.org/10.1016/j.jbiotec.2020.11.004>
- Kumaraswamy RV, Saharan V, Kumari S, Choudhary RC, Pal A, Sharma SS, Rakshit S, Raliya R, Biswas P (2021). Chitosan-silicon nanofertilizer to enhance



- plant growth and yield in maize (*Zea mays* L.). *Plant Physiology and Biochemistry* 159, 53–66. <https://doi.org/10.1016/j.plaphy.2020.11.054>
- Lassalle G (2021). Monitoring natural and anthropogenic plant stressors by hyperspectral remote sensing: Recommendations and guidelines based on a meta-review. *Science of the Total Environment* 788, 147758. <https://doi.org/10.1016/j.scitotenv.2021.147758>
- Lassalle G, Fabre S, Credoz A, Dubucq D, Elger A (2020). Monitoring oil contamination in vegetated areas with optical remote sensing: a comprehensive review. *J. Hazard. Mater.* 393, 122427. <https://doi.org/10.1016/j.jhazmat.2020.122427>
- Li M, Xu G, Huang F, Hou S, Liu B, Yu Y (2021). Influence of nano CuO on uptake and translocation of bifenthrin in rape (*Brassica napus* L.). *Food Control* 130, 108333. <https://doi.org/10.1016/j.foodcont.2021.108333>
- Li T, Xu H (2020). Selenium-Containing Nanomaterials for Cancer Treatment. *Cell Reports Physical Science* 1, 100111. <https://doi.org/10.1016/j.xcrp.2020.100111>
- Lian M, Wang L, Feng Q, Niu L, Zhao Z, Wang P, Song C, Li X, Zhang Z (2021). Thiol-functionalized nano-silica for in-situ remediation of Pb, Cd, Cu contaminated soils and improving soil environment. *Environmental Pollution* 280, 116879. <https://doi.org/10.1016/j.envpol.2021.116879>
- Macůrkova M, L. Maryska L, Jindrichova B, Drobnikova T, Vrchotova B, Pospichalova R, Zaruba K, Hubacek T, Siegel J, Burketova L, Loveck P, Valentova O (2021). Effect of round-shaped silver nanoparticles on the genetic and functional diversity of soil microbial community in soil and “soil-plant” systems. *Applied Soil Ecology* 168, 104165. <https://doi.org/10.1016/j.apsoil.2021.104165>
- Malejko J, Godlewska-Zytkiewicz B, Vanek T, Landa P, Nath J, Dror I, Berkowitz B (2021). Uptake, translocation, weathering and speciation of gold nanoparticles in potato, radish, carrot and lettuce crops. *Journal of Hazardous Materials* 418, 126219. <https://doi.org/10.1016/j.jhazmat.2021.126219>
- Mathur P, Roy S (2020). Nanosilica facilitates silica uptake, growth and stress tolerance in plants. *Plant Physiology and Biochemistry* 157, 114–127. <https://doi.org/10.1016/j.plaphy.2020.10.011>
- Mehta MR, Mahajan HP, Hivrale AU (2021). Green Synthesis of Chitosan Capped-Copper Nano Biocomposites: Synthesis, Characterization, and Biological Activity against Plant Pathogens. *BioNanoScience* 11, 417–427. <https://doi.org/10.1007/s12668-021-00823-8>
- Mishra P, Xue Y, Eivazi F, Afrasiabi Z (2021). Size, concentration, coating, and exposure time effects of silver nanoparticles on the activities of selected soil enzymes. *Geoderma* 381, 114682. <https://doi.org/10.1016/j.geoderma.2020.114682>
- Moradbeygi H, Jamei R, Heidari R, Darvishzadeh R (2020). Investigating the enzymatic and non-enzymatic antioxidant defense by applying iron oxide nanoparticles in *Dracocephalum moldavica* L. plant under salinity stress. *Scientia Horticulturae* 272, 109537. <https://doi.org/10.1016/j.scienta.2020.109537>
- Morales-Espinoza MC, Cadenas-Pliego G, Pérez-Alvarez M, Hernández-Fuentes AD, de la Fuente MC, Benavides-Mendoza A, Valdés-Reyna J, Juárez-Maldonado A (2019). Se Nanoparticles Induce Changes in the Growth, Antioxidant Responses, and Fruit Quality of Tomato Developed under NaCl Stress. *Molecules*, 24, 3030. doi:10.3390/molecules24173030
- Neysanian M, Iranbakhsh A, Ahmadvand R, Oraghi Ardebili Z, Ebadi M (2020). Comparative efficacy of selenate and selenium nanoparticles for improving growth, productivity, fruit quality, and postharvest longevity through modifying nutrition, metabolism, and gene expression in tomato; potential benefits and risk assessment. *PLoS ONE* 15(12), e0244207. <https://doi.org/10.1371/journal.pone.0244207>
- Noman M, Shahid M, Ahmed T, Tahir M, Naqqash T, Muhammad S, Song F, Abid HMA, Aslam Z (2020). Green copper nanoparticles from a native *Klebsiella pneumoniae* strain alleviated oxidative stress impairment of wheat plants by reducing the chromium bioavailability and increasing the growth. *Ecotoxicology and Environmental Safety* 192, 110303. <https://doi.org/10.1016/j.ecoenv.2020.110303>
- Ogunkunle CO, Hussein SO, Okoro K, Adimula V (2021). Interaction of nanoparticles with soil. In: Abdeltif Amrane et al. (Eds.), *Nanomaterials for Soil Remediation*. Elsevier Inc., Pp: 101-132. <https://doi.org/10.1016/B978-0-12-822891-3.00006-2>
- Ogunkunle CO, Odulaja DA, Akande FO, Varun M, Vishwakarma V, Fatoba PO (2020). Cadmium

- toxicity in cowpea plant: Effect of foliar intervention of nano-TiO<sub>2</sub> on tissue Cd bioaccumulation, stress enzymes and potential dietary health risk. *Journal of Biotechnology* 310, 54–61. <https://doi.org/10.1016/j.jbiotec.2020.01.009>
- Othman R, Mohd Zaifuddin FA, Hassan NM (2014). Carotenoid biosynthesis regulatory mechanisms in plants. *J. Oleo Sci.* 63, 753–760. <https://doi.org/10.5650/jos.ess13183>
- Pagano L, Pasquali F, Majumdar S, Torre-Roche RD, Zuverza-Mena N, Villani M, Zappettini A, Marra RE, Isch SM, Marmiroli M, Maestri E, Dhankher OP, White JC, Marmiroli N (2017). Exposure of *Cucurbita pepo* to binary combinations of engineered nanomaterials: physiological and molecular response. *Environ. Sci. Nano* 4, 1579–1590. <https://doi.org/10.1039/C7EN00219J>
- Parveen A, Siddiqui ZA (2021). Impact of silicon dioxide nanoparticles on growth, photosynthetic pigments, proline, activities of defense enzymes and some bacterial and fungal pathogens of tomato. *Vegetos*, <https://doi.org/10.1007/s42535-021-00280-4>
- Paul A, Roychoudhury (2021). Go green to protect plants: repurposing the antimicrobial activity of biosynthesized silver nanoparticles to combat phytopathogens. *Nanotechnology for Environmental Engineering* 6, 10. <https://doi.org/10.1007/s41204-021-00103-6>
- Prakash V, Peralta-Videa J, Tripathi DK, Ma X, Sharma S (2021). Recent insights into the impact, fate and transport of cerium oxide nanoparticles in the plant-soil continuum. *Ecotoxicology and Environmental Safety* 221, 112403. <https://doi.org/10.1016/j.ecoenv.2021.112403>
- Pu S, Yan C, Huang H, Liu S, Deng D (2019). Toxicity of nano-CuO particles to maize and microbial community largely depends on its bioavailable fractions. *Environmental Pollution* 255, 113248. <https://doi.org/10.1016/j.envpol.2019.113248>
- Qi W-Y, Li Q, Chen H, Liu J, Xing S-F, Xu M, Yan Z, Song C, Wang S-G (2021). Selenium nanoparticles ameliorate *Brassica napus* L. cadmium toxicity by inhibiting the respiratory burst and scavenging reactive oxygen species. *Journal of Hazardous Materials* 417, 125900. <https://doi.org/10.1016/j.jhazmat.2021.125900>
- Quiterio-Gutiérrez T, Ortega-Ortiz H, Cadenas-Pliego G, Hernández-Fuentes AD, Sandoval-Rangel A, Benavides-Mendoza A, Cabrera-de la Fuente M, Juárez-Maldonado A (2019). The Application of Selenium and Copper Nanoparticles Modifies the Biochemical Responses of Tomato Plants under Stress by *Alternaria solani*. *Int. J. Mol. Sci.*, 20, 1950. doi:10.3390/ijms20081950
- Rady MM, Desoky E-SM, Ahmed SM, Majrashi A, Ali EF, Arnaout SMAI, Selem E (2021). Foliar Nourishment with Nano-Selenium Dioxide Promotes Physiology, Biochemistry, Antioxidant Defenses, and Salt Tolerance in *Phaseolus vulgaris*. *Plants*, 10, 1189. <https://doi.org/10.3390/plants10061189>
- Rai-Kalal P, Tomar RS, Jajoo A (2021). H<sub>2</sub>O<sub>2</sub> signaling regulates seed germination in ZnO nanoprimed wheat (*Triticum aestivum* L.) seeds for improving plant performance under drought stress *Environmental and Experimental Botany* 189, 104561. <https://doi.org/10.1016/j.envexpbot.2021.104561>
- Rajput V, Minkina T, Mazarji M, Shende S, Sushkova S, Mandzhieva S, Burachevskaya M, Chaplygin V, Singh A, Jatav H (2020). Accumulation of nanoparticles in the soil-plant systems and their effects on human health. *Annals of Agricultural Sciences* 65, 137–143. <https://doi.org/10.1016/j.aas.2020.08.001>
- Rawat S, Adisa IO, Wang Y, Sun Y, Fadil AS, Niu G, Sharma N, Hernandez-Viezcas JA, Peralta-Videa JR, Gardea-Torresdey JL (2019). Differential physiological and biochemical impacts of nano vs micron Cu at two phenological growth stages in bell pepper (*Capsicum annuum*) plant. *NanoImpact* 14, 100161. <https://doi.org/10.1016/j.impact.2019.100161>
- Rizwan M, Ali S, Zia ur Rehman M, Riaz M, Adrees M, Hussain A, Zahir ZA, Rinklebe J (2021). Effects of nanoparticles on trace element uptake and toxicity in plants: A review. *Ecotoxicology and Environmental Safety* 221, 112437. <https://doi.org/10.1016/j.ecoenv.2021.112437>
- Saini S, Kaur N, Pati PK (2021). Phytohormones: Key players in the modulation of heavy metal stress tolerance in plants. *Ecotoxicology and Environmental Safety* 223, 112578. <https://doi.org/10.1016/j.ecoenv.2021.112578>
- Sathiyabama M, Indhumathi M, Amutha T (2020). Preparation and characterization of curcumin functionalized copper nanoparticles and their application enhances disease resistance in chickpea against wilt pathogen. *Biocatalysis and*

- Agricultural Biotechnology 29, 101823. <https://doi.org/10.1016/j.bcab.2020.101823>
- Seliem MK, Hafez Y, El-Ramady H (2020). Using of nano - selenium in reducing the negative effects of high temperature stress on *Chrysanthemum morifolium* Ramat. J. Sustain. Agric. Sci. 46 (3), 47-59. <https://doi.org/10.21608/jsas.2020.23905.1203>
- Servin AD, Pagano L, Castillo-Michel H, De la Torre-Roche R, Hawthorne J, Hernandez-Viezcas JA, Loredó-Portales R, Majumdar S, Gardea-Torresday J, Dhankher OP, White JC (2017). Weathering in soil increases nanoparticle CuO bioaccumulation within a terrestrial food chain. Nanotoxicology 11, 98–111.
- Shah T, Latif S, Saeed F, Ali I, Ullah S, Alsahli AA, Jan S, Ahmad P (2021). Seed priming with titanium dioxide nanoparticles enhances seed vigor, leaf water status, and antioxidant enzyme activities in maize (*Zea mays* L.) under salinity stress. Journal of King Saud University – Science 33, 101207. <https://doi.org/10.1016/j.jksus.2020.10.004>
- Shalaby TA, Abd-Alkarim E, El-Aidy F, Hamed E, Sharaf-Eldin M, Taha N, El-Ramady H, Bayoumi Y, Reis ARD (2021). Nano-selenium, silicon and H<sub>2</sub>O<sub>2</sub> boost growth and productivity of cucumber under combined salinity and heat stress. Ecotoxicology and Environmental Safety 212, 111962. <https://doi.org/10.1016/j.ecoenv.2021.111962>
- Sharifan H, Moore J, Ma X (2020). Zinc oxide (ZnO) nanoparticles elevated iron and copper contents and mitigated the bioavailability of lead and cadmium in different leafy greens. Ecotoxicol. Environ. Saf. 191, 110177 <https://doi.org/10.1016/j.ecoenv.2020.110177>
- Sharma S, Kumar K, Thakur N, Chauhan MS (2020). *Ocimum tenuiflorum* leaf extract as a green mediator for the synthesis of ZnO nanocapsules inactivating bacterial pathogens. Chemical Papers 74, 3431–3444. <https://doi.org/10.1007/s11696-020-01177-3>
- Sheikhalipour M, Esmailpour B, Behnamian M, Gohari G, Giglou MT, Vachova P, Rastogi A, Brestic M, Skalicky M (2021). Chitosan–Selenium Nanoparticle (Cs–Se NP) Foliar Spray Alleviates Salt Stress in Bitter Melon. Nanomaterials (Basel) 11(3), 684. doi: 10.3390/nano11030684
- Shrivastava M, Srivastav A, Gandhi S, Rao S, Roychoudhury A, Kumar A, Singhal RK, Jha SK, Singh SD (2019). Monitoring of engineered nanoparticles in soil-plant system: A review. Environmental Nanotechnology, Monitoring & Management 11, 100218. <https://doi.org/10.1016/j.enmm.2019.100218>
- Simonin M, Martins JMF, Uzu G, Spadini L, Navel A, Richaume A (2021). Low mobility of CuO and TiO<sub>2</sub> nanoparticles in agricultural soils of contrasting texture and organic matter content. Science of the Total Environment 783, 146952. <https://doi.org/10.1016/j.scitotenv.2021.146952>
- Singh D, Kumar A (2020). Quantification of metal uptake in *Spinacia oleracea* irrigated with water containing a mixture of CuO and ZnO nanoparticles. Chemosphere 243, 125239. <https://doi.org/10.1016/j.chemosphere.2019.125239>
- Sreelakshmi B, Induja S, Adarsh PP, Rahul HL, Arya SM, Aswana S, Haripriya R, Aswathy BR, Manoj PK, Vishnudasan D (2021). Drought stress amelioration in plants using green synthesised iron oxide nanoparticles. Materials Today: Proceedings 41, 723–727. <https://doi.org/10.1016/j.matpr.2020.05.801>
- Tauseef A, Hisamuddin, Khalilullah A, Uddin I (2021). Role of MgO nanoparticles in the suppression of *Meloidogyne incognita*, infecting cowpea and improvement in plant growth and physiology. Experimental Parasitology 220, 108045. <https://doi.org/10.1016/j.exppara.2020.108045>
- Tavallali V, Rowshan V, Gholami H, Hojati S (2020). Iron-urea nano-complex improves bioactive compounds in essential oils of *Ocimum basilicum* L. Scientia Horticulturae 265, 109222. <https://doi.org/10.1016/j.scienta.2020.109222>
- Wang C, Cheng T, Liu H, Zhou F, Zhang J, Zhang M, Liu X, Shi W, Cao T (2021a). Nano-selenium controlled cadmium accumulation and improved photosynthesis in indica rice cultivated in lead and cadmium combined paddy soils. J Environ sci 103, 336–346. <https://doi.org/10.1016/j.jes.2020.11.005>
- Wang CR, Rong H, Zhang XB, Shi WJ, Hong X, Liu WC (2020a). Effects and mechanisms of foliar application of silicon and selenium composite sols on diminishing cadmium and lead translocation and affiliated physiological and biochemical responses in hybrid rice (*Oryza sativa* L.) exposed to cadmium and lead. Chemosphere, 251, Article 126347. <https://doi.org/10.1016/j.chemosphere.2020.126347>
- Wang D, Rensing C, Zheng S (2022). Microbial reduction and resistance to selenium: Mechanisms, applications and prospects. Journal of Hazardous Materials 421, 126684. <https://doi.org/10.1016/j.jhazmat.2021.126684>

- Wang K, Wang Y, Li K, Wan Y, Wang Q, Zhuang Z, Guo Y, Li H (2020b). Uptake, translocation and biotransformation of selenium nanoparticles in rice seedlings (*Oryza sativa* L.). *J Nanobiotechnol* 18, 103. <https://doi.org/10.1186/s12951-020-00659-6>
- Wang Z, Yue L, Dhankher OP, Xing B (2020c). Nano-enabled improvements of growth and nutritional quality in food plants driven by rhizosphere processes. *Environ. Int.* 142, 105831 <https://doi.org/10.1016/j.envint.2020.105831>.
- Ye W, Yuan L, Zhu R, Yin X, Banuelos G (2021). Selenium volatilization from tundra soils in maritime Antarctica. *Environment International* 146, 106189. <https://doi.org/10.1016/j.envint.2020.106189>
- Zahedi SM, Moharrami F, Sarikhani S, Padervand M (2020). Selenium and silica nanostructure-based recovery of strawberry plants subjected to drought stress. *Sci Rep.* 10: 17672. doi: 10.1038/s41598-020-74273-9
- Zhang J-Y, Zhou H, Zeng P, Wang S-L, Yang W-J, Huang F, Huo Y, Yu S-N, Gu J-F, Liao B-H (2021). Nano-Fe<sub>3</sub>O<sub>4</sub>-modified biochar promotes the formation of iron plaque and cadmium immobilization in rice root. *Chemosphere* 276, 130212. <https://doi.org/10.1016/j.chemosphere.2021.130212>